Supplementary Material for:

Antitumor Activities and Cellular Changes Induced by TrkB Inhibition in Medulloblastoma

Amanda Thomaz ^{1,2} • Kelly de Vargas Pinheiro ^{1,2} • Bárbara Kunzler Souza ^{1,2} • Lauro Gregianin ^{1,4,5} • Algemir L. Brunetto ^{1,3} • André T. Brunetto ^{1,3} • Caroline Brunetto de Farias ^{1,3} • Mariane da Cunha Jaeger ^{1,3} • Vijay Ramaswamy ^{6,7} • Carolina Nör ^{6,8} • Michael D. Taylor ^{6,7,8,9,10} • Rafael Roesler ^{1,2}*

¹Cancer and Neurobiology Laboratory, Experimental Research Center, Clinical Hospital (CPE-HCPA), Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil, ² Department of Pharmacology, Institute for Basic Health Sciences, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil, ³ Children's Cancer Institute, Porto Alegre, RS, Brazil, ⁴ Department of Pediatrics, School of Medicine, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil, ⁵ Pediatric Oncology Service, Clinical Hospital, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil, ⁶ The Arthur and Sonia Labatt Brain Tumour Research Centre, The Hospital for Sick Children, Toronto, ON, Canada, ⁷ Division of Haematology/Oncology, The Hospital for Sick Children, Toronto, ON, Canada, ⁸ Developmental and Stem Cell Biology Program, The Hospital for Sick Children, Toronto, ON, Canada, ⁹ Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada, ¹⁰ Division of Neurosurgery, The Hospital for Sick Children, Toronto, ON, Canada.

* Correspondence:


Rafael Roesler, Department of Pharmacology, Institute for Basic Health Sciences, Federal


University of Rio Grande do Sul, Rua Sarmento Leite, 500 (ICBS, Campus Centro/UFRGS), 90050-170 Porto Alegre, RS, Brazil.


Telephone: +5551 33083183; fax: +5551 33083121.


E-mail: rafaelroesler@hcpa.edu.br


SUPPLEMENTARY FIGURE S1 Original Western blots membranes analysed by ImageQuant LAS500 (GE Healthcare Life Sciences, Little Chalfont, UK) and original membranes stained with Coomassie blue (0.025%) for western blots in Figures 5 and 6. Membranes were cut prior to antibody stainings to allow for detection of proteins running at different sizes on the same membrane.

















ANA-12 [μM] 24 h

